

BRAZILIAN BUTT LIFT

Pre-Op Instructions for Brazilian Butt Lift (BBL)

Congratulations on taking the first step towards achieving your aesthetic goals with your Brazilian Butt Lift surgery. To ensure a smooth and successful procedure, please carefully follow the pre-operative instructions outlined below:

Night Before Surgery:

- **Nothing to Eat or Drink:** Please refrain from eating or drinking anything within 8 hours of your surgery, not even water. Failure to comply with this instruction may result in the cancellation of your surgery. However, you may take certain medications with a small sip of water (We will review these medications with you).
- **Antibiotics:** Take your first dose of antibiotics the night before surgery.
- **Set Up Home Recovery Area:** Prepare a comfortable recovery area in your home. This may include pillows, blankets, puppy pads, books, television, and any other items that will help make your recovery as comfortable as possible.
- **Bathing:** The night before surgery you may take a shower with antibacterial soap, but please avoid applying lotion, perfume or other skincare products.

Day of Surgery:

- **Dress Comfortably:** Wear comfortable, clean, and loose-fitting clothes on the day of your surgery. We recommend a warm sleeveless, loose-fitting shirt that can be buttoned or zipped, or robe and slippers.
- **Avoid Make up, Jewelry, and Nails:** Please do not wear any makeup, jewelry, cosmetic creams, deodorant, or piercings on the day of your surgery. Avoid long, thick acrylic nails.
- **Medications:** Take any medications that our provider has told you to take the morning of surgery with only a sip of water. Bring your medications with you on surgery day.
- **Transportation:** Make sure you have arranged for someone to pick you up after surgery. You will provide us with their contact information. If you receive general anesthesia, you will not be able to drive yourself or take a taxi/uber. You may bring your BBL pillow for the ride home.

What to bring with you:

- **Pre-op folder:** Bring your folder with all of your post op instructions
- **Medications:** Bring all prescriptions with you on surgery day.
- **Compression socks/TED hose:** Put your compression socks on prior to surgery.
- **Backpack:** Bring a back pack or small bag to keep your belongings in. Leave valuables at home unless you have a payment that is due the day of surgery.
- **Blanket:** Bring a small blanket.

By following these pre-operative instructions diligently, you will help ensure the safety and success of your BBL procedure. If you have any questions or concerns, please do not hesitate to contact our office at **214-227-0668**.

We look forward to assisting you on your journey to a more confident and rejuvenated you.

Sincerely,
Noble Cosmetic Surgery

Post-Operative Instructions for Brazilian Butt Lift (BBL)

It is essential to follow these post-operative instructions carefully to ensure a smooth and comfortable healing process:

Medications:

- **Pain Management:** You will be prescribed pain medication to alleviate post-operative discomfort. If your pain is mild, you may opt to take Tylenol instead, but avoid combining it with the prescribed medication. Take your pain medication with food to minimize the risk of nausea. Do not take any NSAIDS (Ex: Aspirin, Ibuprofen, Motrin, Excedrin)
- **Antibiotics:** Take your antibiotics as prescribed. Finish all antibiotics.
- **Nausea:** Nausea is common after surgery, especially with anesthesia and strong medications. If you experience nausea, take the prescribed medication as directed.
- **Arnica:** Take over the counter arnica tablets as directed. This will help with bruising and swelling.
- **Constipation:** Narcotic pain relievers and anesthesia can cause constipation. Start taking an over-the-counter stool softener (Dulcolax soft chews, Colace, Miralax) to avoid constipation.
- **Medications to Avoid:** Only take medications approved or prescribed by your surgeon to prevent complications.

Nutrition:

- **Diet:** Begin with a light, low-fat diet after surgery. You can transition to a regular diet as long as you don't experience nausea or vomiting. Start with liquids and gradually progress to solid foods. Watch your sodium intake. Salty foods will cause you to swell more. Increase your protein intake, which will help with wound healing.
- **Hydration:** Drink 8-10 glasses of water daily to stay hydrated. Even if you don't have much of an appetite make sure to drink plenty of water and protein shakes. Avoid alcohol for at least 48 hours and while taking pain medications.

Activity:

- **Sitting:** Do not sit directly on your buttocks for 3 weeks. Whenever you must sit, use your BBL pillow.
- **Movement:** It's crucial to move around immediately after surgery. Move your legs and ankles while lying down and take deep breaths to keep your lungs clear. This decreases your chances of developing blood clots and pneumonia. Make sure to get up and walk every 2 hours while awake.

- **Exercise:** Start off with walking and low impact exercises as soon as you feel up to it. Any strenuous exercise or activity will increase your blood flow and cause more swelling, so gradually work back up to your regular routine.
- **Compression Garment:** Wear the provided compression garment for 6 weeks to minimize swelling and aid in body contouring. Ensure it fits snugly but not too tight. We will fit you in your compression garment at your follow up appt.
- **Driving:** Refrain from driving or making important decisions while on pain medications. Wait at least 6 hours after discontinuing pain medication before operating a vehicle.
- **Travel:** Take breaks during automobile travel every 2 hours to prevent blood pooling and clots. Avoid airline travel for the first week post-surgery due to increased swelling.
- **Return to Work:** Most patients require anywhere from 3 to 5 days off work, depending on job responsibilities. You will need to be standing at work or using a BBL pillow when sitting.
- **Sexual Intercourse:** You may resume sexual activity once your incisions are fully healed. Do not place any pressure on your buttocks for 3 weeks
- **Sleeping:** You will sleep on you stomach or on your back with pillows to prevent you from placing any pressure on your buttocks for 3 weeks.

Bathing:

- **Showering:** Typically, you can shower the day after surgery depending on what other surgeries you had. Check with the doctor for instructions on whether you can shower. You may sponge bath until cleared. Do not use any scar treatment, ointments, lotions, or other skincare creams on the incision until instructed to do so.
- **Hot Tubs/Baths/Swimming Pools:** No hot tubs, baths, or swimming pools until cleared by the doctor.

Please adhere to these instructions closely to optimize your recovery and achieve the best possible results. If you have any questions or concerns, do not hesitate to contact our office.

Wishing you a speedy and comfortable recovery!

Post-Care Instructions: Incisions

How to Take Care of Your Incisions:

- **Incisions:** You will have drainage for approximately 2-3 days from your liposuction incisions. You will change the maxi pads as needed whenever they are soaked. Do not place bandaids over the incisions. Do not use any scar treatment, ointments, lotions, or other skincare creams directly on the incision until fully healed with no scabs.
- **Stitches:** You will have some stitches around your buttocks from the fat transfer. We will remove these at your follow up.

What to Expect:

- **Drainage:** Drainage from incision sites is normal for the first 24-72 hours, usually blood-tinged. If needed, use gauze or a light pad to reinforce post-op dressings.
- **Bruising:** Expect bruising, which can be significant depending on the extent of liposuction. Bruises typically resolve within 2-3 weeks, changing from purple to yellow/green as they heal. You may take arnica to help with bruising.
- **Swelling:** Swelling is common for weeks to months after liposuction. Arnica, compression garments, and rest can help manage swelling.
- **Itching:** Itching around incision sites is normal and can be relieved with Benadryl or a non-drowsy antihistamine if necessary.
- **Sensory Changes:** Numbness, tingling, pulling/pinching, sharp shooting pain around incisions and liposuction areas. This is normal and should improve over time.
- **Scarring:** Scars are minimal and may take up to a year to fully heal. Tape or topical silicone gel can aid in scar management. Do not use any scar treatment on incisions until cleared by the doctor.

Precautions - What Not to Do:

- **Avoid Soaking:** Do not soak in baths, jacuzzi, or hot tubs until cleared by the doctor.
- **Avoid Blood Thinners:** Refrain from taking aspirin, ibuprofen, naproxen, or other blood thinners until cleared by your surgeon.
- **No Heating Pads or Ice Packs:** Unless instructed otherwise by your surgeon, do not apply heating pads or ice packs to treated areas.

Emergency Situations - When to Seek Medical Attention:

- **Signs of Infection:** Including spreading redness, increased drainage or pus, worsening pain, warmth at incision site, and fever over 101°F.
- **Excessive Bleeding:** If dressings are saturated with bright red blood and require frequent changes.
- **Other Emergencies:** Such as difficulty breathing, chest pain, severe vomiting, or asymmetric swelling in your legs.

For any concerns, please call our office at 214-227-0668 or seek immediate medical attention by calling 911.

We're here to support you throughout your recovery journey.